

# 1-(9*H*-Carbazol-4-yloxy)-3-(2-(2-[<sup>11</sup>C]methoxyphenoxy)ethylamino)-propan-2-ol [<sup>11</sup>C]Carvedilol

Created: December 30, 2005

Updated: January 17, 2006

**Chemical name:** 1-(9*H*-Carbazol-4-yloxy)-3-(2-(2-[<sup>11</sup>C]methoxyphenoxy)ethylamino)-propan-2-ol

**Abbreviated name:** [<sup>11</sup>C]CARV

**Synonym:** [<sup>11</sup>C]Carvedilol

**Backbone:** Compound

**Target:** P-glycoprotein multidrug transporter, MDR-1

**Mechanism:** P-glycoprotein substrate

**Method of detection:** PET

**Source of signal:** <sup>11</sup>C


**Activation:** No

***In vitro* studies:** Yes

**Rodent studies:** Yes

**Other non-primate mammal studies:** No

**Non-human primate studies:** No

**Human studies:** No

Click on the above structure for additional information in PubChem  
[<http://pubchem.ncbi.nlm.nih.gov>].

## Background

[PubMed]

One of the mechanisms cells use to escape the cytotoxic effects of chemotherapeutic agents, such as Adriamycin, Vinca alkaloids, epipodophyllotoxins, actinomycin D, and Taxol<sup>®</sup>, is to limit their presence inside the cells through the actions of P-glycoprotein (P-gp), a protein encoded by the multidrug resistance (MDR-1) gene (1, 2). P-gp is an ATP-dependent transmembrane multidrug transporter that is capable of actively pumping a variety of agents out of cells. Overexpression of P-gp in tumor cells (such as renal carcinoma, hepatoma, pheochromocytoma, and colon carcinoma) leads to resistance to anticancer drugs (3). P-gp is also present in a variety of normal cells, such as intestinal mucosal cells, hepatocytes, renal proximal tubule epithelial cells, and endothelial cells of

the blood-brain barrier (4, 5). Calcium channel blockers, cyclosporin A, and its non-immunosuppressive analog PSC 833 are MDR modulators that inhibit transport of P-gp substrates out of cells (6, 7).

Sestamibi (MIBI) is a substrate for P-gp. <sup>99m</sup>Tc-MIBI [<http://www.ncbi.nlm.nih.gov/books/bv.fcgi?rid=micad.chapter.MIBI99mtc>] has been approved by the United States Food and Drug Administration as a myocardial perfusion imaging agent for use in single-photon emission computed tomography (SPECT) to assess the risk of future cardiac events. It is also approved as a tumor-imaging agent in breast, lung, thyroid, and brain cancers. 1-(9*H*-Carbazol-4-yloxy)-3-[2-(2-methoxyphenoxy)ethylamino]-propan-2-ol (carvedilol), a  $\beta$ -adrenoceptor antagonist, is also a transport substrate for P-gp with a log *P* of 2.0 (8). [<sup>11</sup>C]Carvedilol ([<sup>11</sup>C]CARV) was found to be not suitable as a tracer for imaging adrenoceptors in any target organ, such as the brain, heart, lungs, and spleen (9). Brain accumulation of [<sup>11</sup>C]CARV is reported to be highly sensitive to P-gp modulation by cyclosporin A (10). Therefore, [<sup>11</sup>C]CARV is being developed as a positron emission tomography (PET) agent to noninvasively study P-gp function.

## Synthesis

---

[PubMed]

In the report by Bart et al. (10), [<sup>11</sup>C]CARV was synthesized by alkylation of desmethylcarvedilol with [<sup>11</sup>C]methyltriflate, with subsequent separation by high-performance liquid chromatography. The radiochemical yield was 20-33%, based on [<sup>11</sup>C]methyltriflate, with a total synthesis time of 55 min. The specific activity was 13-26 TBq/mmol (350-703 Ci/mmol) at the end of bombardment.

## In Vitro Studies: Testing in Cells and Tissues

---

[PubMed]

In a study using the P-gp-negative small cell lung carcinoma GLC<sub>4</sub> and its overexpressing subclone GLC<sub>4</sub>/P-gp, cellular accumulation of [<sup>11</sup>C]CARV was 100% higher in GLC<sub>4</sub> than in GLC<sub>4</sub>/P-gp (10). Cyclosporin A (50  $\mu$ M) increased [<sup>11</sup>C]CARV accumulation by 2.0-fold (*P* = 0.0001) in GLC<sub>4</sub>/P-gp but had no effect in GLC<sub>4</sub>. MK571 (50  $\mu$ M), a multidrug resistance-associated protein (MRP1) inhibitor, had no effect on [<sup>11</sup>C]CARV accumulation in both cell types.

## Animal Studies

---

### Rodents

[PubMed]

Biodistribution studies in normal rats showed high accumulation of radioactivity, as measured by autoradiography, in the lung, followed by the liver, kidney, and heart at 60 min after injection of [<sup>11</sup>C]CARV (10). Tracer accumulation in the brain was low. Cyclosporin A (25-50 mg/kg) increased [<sup>11</sup>C]CARV accumulation in the brain 6  $\pm$  3-fold. This indicates that P-gp-mediated [<sup>11</sup>C]CARV efflux

can be modulated by cyclosporin A. PET imaging revealed that [<sup>11</sup>C]CARV accumulated in the brains of rats without or with cyclosporin A treatment (5, 10, 15, 25, and 50 mg/kg). The tracer was rapidly cleared from plasma after injection, with a first phase half-life of 2.3 min and a second, wash-out phase half-life of 108 min. Cyclosporin A had no effects on the plasma kinetics of [<sup>11</sup>C]CARV. The distribution volume (DV) of [<sup>11</sup>C]CARV in the brain was 0.3 after injection of 35-50 MBq (0.94-1.35 mCi) of [<sup>11</sup>C]CARV. At 5 mg/kg cyclosporin A, the DV was increased >1-fold. The maximum increase in DV was 2-fold after 15 mg/kg cyclosporin A with no further increases at higher dosages of cyclosporin A. Metabolism studies using [<sup>14</sup>C]CARV in rats, mice, and dogs showed that all metabolites are more hydrophilic than the parent compound and are expected to have negligible brain uptake (11).

## Other Non-Primate Mammals

[PubMed]

No relevant publications are currently available.

## Non-Human Primates

[PubMed]

No publications are currently available.

## Human Studies

---

[PubMed]

No relevant publications are currently available.

## References

---

1. Endicott JA, Ling V. The biochemistry of P-glycoprotein-mediated multidrug resistance. *Annu Rev Biochem* 58:137–171; 1989. (PubMed)
2. Gottesman MM, Pastan I. Biochemistry of multidrug resistance mediated by the multidrug transporter. *Annu Rev Biochem* 62:385–427; 1993. (PubMed)
3. Fojo AT, Ueda K, Slamon DJ, Poplack DG, Gottesman MM, Pastan I. Expression of a multidrug-resistance gene in human tumors and tissues. *Proc Natl Acad Sci U S A* 84(1):265–269; 1987. (PubMed)
4. Piwnicka-Worms D, Rao VV, Kronauge JF, Croop JM. Characterization of multidrug resistance P-glycoprotein transport function with an organotechnetium cation. *Biochemistry* 34(38):12210–12220; 1995. (PubMed)
5. Thiebaut F, Tsuruo T, Hamada H, Gottesman MM, Pastan I, Willingham MC. Cellular localization of the multidrug-resistance gene product P-glycoprotein in normal human tissues. *Proc Natl Acad Sci U S A* 84(21):7735–7738; 1987. (PubMed)
6. Hughes CS, Vaden SL, Manaugh CA, Price GS, Hudson LC. Modulation of doxorubicin concentration by cyclosporin A in brain and testicular barrier tissues expressing P-glycoprotein in rats. *J Neurooncol* 37(1): 45–54; 1998. (PubMed)
7. Mayer U, Wagenaar E, Dorobek B, Beijnen JH, Borst P, Schinkel AH. Full blockade of intestinal P-glycoprotein and extensive inhibition of blood-brain barrier P-glycoprotein by oral treatment of mice with PSC833. *J Clin Invest* 100(10):2430–2436; 1997. (PubMed)

8. Jonsson O, Behnam-Motlagh P, Persson M, Henriksson R, Grankvist K. Increase in doxorubicin cytotoxicity by carvedilol inhibition of P-glycoprotein activity. *Biochem Pharmacol* 58(11):1801–1806; 1999. (PubMed)
9. Doze P, Elsinga PH, Maas B, Van Waarde A, Wegman T, Vaalburg W. Synthesis and evaluation of radiolabeled antagonists for imaging of beta-adrenoceptors in the brain with PET. *Neurochem Int* 40(2): 145–155; 2002. (PubMed)
10. Bart J, Dijkers EC, Wegman TD, de Vries EG, van der Graaf WT, Groen HJ, Vaalburg W, Willemsen AT, Hendrikse NH. New positron emission tomography tracer [(11)C]carvedilol reveals P-glycoprotein modulation kinetics. *Br J Pharmacol* 145(8):1045–1051; 2005. (PubMed)
11. Schaefer WH, Politowski J, Hwang B, Dixon F, Goalwin A, Gutzait L, Anderson K, DeBrosse C, Bean M, Rhodes GR. Metabolism of carvedilol in dogs, rats, and mice. *Drug Metab Dispos* 26(10):958–969; 1998. (PubMed)